

Article ID: 195528  
DOI: 10.5586/am/195528

**Publication History**  
Received: 2024-07-10  
Accepted: 2024-11-03  
Published: 2024-12-03

**Handling Editor**  
Piotr Zaniewski; Warsaw  
University of Life Sciences,  
Warsaw, Poland;  
<https://orcid.org/0000-0002-0792-9854>


**Funding**  
Research financed from  
statutory funds of the Faculty of  
Biology and Biotechnology,  
UWM in Olsztyn.

**Competing Interests**  
No competing interests have  
been declared.

**Copyright Notice**  
© The Author(s) 2024. This is an  
open access article distributed  
under the terms of the [Creative  
Commons Attribution License](#),  
which permits redistribution,  
commercial and  
noncommercial, provided that  
the article is properly cited.

## ORIGINAL RESEARCH

# Problems and challenges of lichen inventory in a changing environment: The case of the “Mszar” and “Redykajny” reserves in Olsztyn

Dariusz Kubiak \*

Department of Microbiology and Mycology, Faculty of Biology and Biotechnology, University of Warmia and Mazury in Olsztyn, Oczapowskiego 1A, 10-719 Olsztyn, Poland

\* Email: [darkub@uwm.edu.pl](mailto:darkub@uwm.edu.pl)

## Abstract

The paper presents the state of knowledge of the lichen biota of the “Mszar” and “Redykajny” nature reserves, located in the “Las Miejski” urban forest complex in Olsztyn (north-eastern Poland). The results of the inventory conducted in 2024 are described, which are compared with published data collected in 1999–2001. So far, a total of 118 lichenized and three non-lichenized fungi have been identified in this area. Recent studies did not confirm the occurrence of 18 species, but at the same time showed 32 species new to the study area. The article discusses possible causes of changes in the species composition of the lichen biota over almost 25 years and the effectiveness of various methods of collecting data in the field. The results prove that both reserves are still important refuges of forest lichen biodiversity on a scale larger than just the area of Olsztyn’s urban forests.

## Keywords

lichenized fungi; species diversity; protected area; Poland

## 1. Introduction

Lichens are an important component of forest communities, constituting a significant part of their biodiversity (Ellis, 2012; Faliński & Mułenko, 1996; Sillet & Antoine, 2004). The most characteristic group of forest lichens are epiphytes, which exclusively, or at least primarily, inhabit the vast space created by trunks and crowns of living trees (Cieśliński et al., 1996; Ellis, 2012). Many lichen species also inhabit dead wood, as facultative or obligate epixylites (Chlebicki et al., 1996; Gutowski et al., 2023). The occurrence of lichen species and the diversity of the communities they create depend on many factors, the most important of which are the presence of suitable substrates and microhabitats (Cieśliński et al., 1996; Sillet & Antoine, 2004) and the specific forest microclimate (Gauslaa, 2014; Phinney et al., 2018). It is known that natural forest communities, usually with a mixed tree species composition and diverse age structure (Peterken, 1996), contain a much larger number of species than commercial forests, which are poorer in microhabitats and substrates (Bergamini et al., 2005; Boch et al., 2013; Hämäläinen et al., 2023; Tripp et al., 2019). Many lichen species have become dependent on these specific conditions and do not occur in managed forests (Bogges et al., 2024; Scheidegger & Stofer, 2015). Most of them are sensitive to changes in environmental conditions (Czerepko et al., 2021), which has been used in

forest bioindication (Fрати & Brunialti, 2023; Miller et al., 2020). Assessment of lichen diversity can also provide reliable information on processes occurring both locally and globally. Many natural and anthropogenic substances are transported over long distances from emission sources and are spread mainly by wet and dry deposition (Fрати & Brunialti, 2023). Lichens have long been used in bioindication and monitoring of transboundary atmospheric pollution (Nash & Gries, 1991; Nimis et al., 2002) and, more recently, global warming (Aptroot et al., 2021; Stanton et al., 2023; Wroblewski et al., 2023). Changes in the species composition of the lichen biota can also inform about potential threats to human health (Cislaghi & Nimis, 1997; Fрати & Brunialti, 2023; Loppi, 2014), therefore it is important to initiate, continue and expand research on lichen diversity in areas located near human settlements. Due to the close proximity to the city and the high risk of exposure to many harmful environmental factors, monitoring the condition of urban forests seems particularly justified.

The aim of the study was to investigate the species diversity of lichen biota in the “Mszar” and “Redykajny” nature reserves in Olsztyn (north-eastern Poland) and to determine possible changes in its composition over the past 25 years.



**Figure 1** Swamp birch forest *Betuletum pubescentis*, “Mszar” nature reserve, locality no. 4 (D. Kubiak).

## 2. Materials and methods

### 2.1. Study site

The “Mszar” and “Redykajny” nature reserves are located within the administrative borders of the city of Olsztyn (over 170,000 inhabitants) in the large (over 1,400 ha) “Las Miejski” woodland complex (Kubiak, 2008). According to the current geographical division of Poland (Solon et al., 2018), both reserves are located in the Olsztyn Lakeland mesoregion. According to the national ATPOL system of grid squares (Verey, 2017), adapted for the plotting of lichen stands (Cieśliński & Fałtynowicz, 1993), both reserves are located in square Be42. The “Redykajny” reserve was established in 1949 over an area of 9.96 ha, while the “Mszar” reserve was established in 1953 over an area of 5.24 ha. The aim of conservation efforts in both reserves is the preservation of raised bogs and boggy forest and shrub communities (CRFOP, 2024; Figure 1, Figure 2, Table 1). Although both areas obtained the status of protected areas in 1907, the first studies of lichen biota were conducted in 1999–2001 (Kubiak, 2008).

### 2.2. Field research

Field data were collected in 2024. Unlike the previous inventory (cf. Kubiak, 2008), information on the occurrence of species and the substrates they inhabited was linked to specific field points. These points were designated for all types of communities in the study area. The number of points in a specific type of community corresponded to its share in the total area of the reserve. Data were collected around a previously designated point at a distance of 10 m (over 300 m<sup>2</sup>) on all substrates inhabited by these organisms. At each point, geograph-

ical coordinates (WGS 1984) were recorded, and the type of local community was determined. The list and general characteristics of the research sites are presented in Table 1. In total, a field study was carried out at 23 sites, including seven within the “Mszar” reserve and 16 within the “Redykajny” reserve. Species that were identified in the field were recorded without collecting specimens. For the remaining taxa, small fragments of thallus were collected for further morphological, anatomical, and chemical studies.

### 2.3. Species identification

Standard spot tests and chromatographic analysis (TLC, solvent C) were used to identify the collected specimens (Orange et al., 2001). The nomenclature of the lichen species follows Fałtynowicz et al. (2024). The categories of lichen threat in Poland are given according to Cieśliński et al. (2006), and the indicator species of lowland primeval forests are given according to Czyżewska and Cieśliński (2003). The collected herbarium material was deposited in the lichen herbarium at the Department of Microbiology and Mycology of the University of Warmia Masuria in Olsztyn (OLTC-L). As part of this part of the study, taxonomic re-identification of specimens collected in 1999–2001 and deposited in OLTC-L was also performed.

## 3. Results and discussion

### 3.1. General characteristics of the lichen biota

As a result of research conducted in 2024, 100 lichenized and three non-lichenized fungi were identified, including 71 in the “Mszar” reserve and 90 in the “Redykajny” reserve (all non-lichenized fungi were recorded in the “Redykajny” reserve).



**Figure 2** Boggy pine forest *Vaccinio uliginosi-Pinetum*, “Redykajny” nature reserve, locality no. 2 (D. Kubiak).

After comparing these results with data from 1999–2001, the total number of species recorded in the study area increased to 121 (Table 2). The latest studies resulted in an increase of this number by 32 species, of which 18 are new to the “Mszar” reserve, and 26 are new to the “Redykajny” reserve. However, the study did not confirm the occurrence of 18 previously recorded species. It should be noted that two species were given in the report of the first inventory (Kubiak, 2008), i.e., *Bacidina assulata* (Körb. S. Ekman and *Placynthiella uliginosa* (Schrad.) Coppins & P. James, are not included in Table 2. The specimen of the first species deposited in OLTC-L and revised in 2024 belongs to *Bacidia arceutina* (Ach.) Arnold) and the second to *P. dasaea* (Stirt.) Tønsberg.

The list of species recorded in both reserves includes 29 threatened and near threatened species which are included in the national Red List, and representing the following categories: 1 – Critically Endangered (*Chaenotheca chlorella*), 8 – Endangered (*Anaptychia ciliaris*, *Bacidia arceutina*, *Calicium trabinellum*, *Cetraria sepincola*, *Chaenotheca brachypoda*, *Ch. stemonea*, *Toniniopsis separabilis*, *Usnea subfloridana*), 10 – Vulnerable (*Bacidia rubella*, *Biatora efflorescens*, *Bryoria fuscescens*, *Chaenotheca xyloxena*, *Nephromopsis chlorophylla*, *Ochrolechia bahusiensis*, *Parmelia submontana*, *Peltigera praetextata*, *Ramalina farinacea*, *Usnea hirta*), and 10 – Near Threatened (*Chaenotheca furfuracea*, *Ch. trichialis*, *Evernia prunastri*, *Graphis scripta*, *Hypogymnia tubulosa*, *Lecanora sarcopisoides*, *Lichenomphalia umbellifera*, *Micarea melaena*, *Pertusaria coccodes*, *Vulpicida pinastri*). The list also includes 13 species legally protected at the national level. Of this group, five species are under strict protection (*A. ciliaris*, *C. sepincola*, *P. submontana*, *P. praetextata*, *Usnea subfloridana*) and eight are under partial protection (*B. fuscescens*, *Cladonia arbuscula*, *H. tubulosa*, *Melanelixia subaurifera*, *N. chlorophylla*, *Ramalina farinacea*, *Usnea hirta*, *V. pinastri*). Four of the

above-mentioned species have the status of lowland primeval forest indicators in Poland: *C. trabinellum*, *Ch. brachypoda*, *Ch. chlorella*, and *M. melaena*.

### 3.2. Lichen species new to the study area

The significant increase in the number of species between the two study periods raises the question of the cause of this change. The most obvious explanation is that the newly discovered species were not previously known and were described as new to science within the last few years (Czarnota & Guzow-Krzemińska, 2010; Guzow-Krzemińska et al., 2016, 2017). However, these species are not numerous (*Lecanora stanislai*, *Micarea byssacea*, *M. soralifera*), and there is a predominance of well-known taxa (both rare and even common) with an established taxonomic position and quite well-understood ecology (e.g. *Chaenotheca chrysocephala*, *Graphis scripta*, *H. tubulosa*, *Lecanora argentata*, *Lepora albescens*, *Physcia adscendens*, *Polycauliona polycarpa*). It may seem that due to the sessile and perennial lifestyle and lack of seasonality in this group of organisms – apart from a few exceptions (Poelt & Vězda, 1990), identification of the majority of lichen species in the field should be simple, compared to some plants or other groups of fungi (von Hirschheydt et al., 2024). Nonetheless, in practice, the number of identified species typically differs from the actual number, and these differences are not limited only to the most inconspicuous, poorly known, or as yet not described taxa. As many field experiments have shown (Vondrák et al., 2016, 2018; von Hirschheydt et al., 2024), the assessment of the biodiversity of lichens may be a challenge even for expert taxonomists. Variables influencing the inventory result can be considered from the perspective of the used research method and environmental heterogeneity. In terms of the process of the inventory (excluding the previ-

**Table 1** Description and geographical coordinates of the study sites in the “Mszar” and “Redykajny” reserves (Poland).

Study sites (no.)	Geographical coordinates (latitude/longitude)	Description (present and potential plant community)
<b>Mszar</b>		
1	53°47'15.8"N, 20°27'51.9"E	Pine-birch swamp forest <i>Thelypteridi-Betuletum pubescentis</i>
2	53°47'17.4"N, 20°27'55.7"E	Alder swamp forest <i>Ribeso nigri-Alnetum</i>
3	53°47'17.5"N, 20°27'53.5"E	Swamp birch forest <i>Betuletum pubescentis</i>
4	53°47'19.2"N, 20°27'54.6"E	Swamp birch forest <i>Betuletum pubescentis</i> (Figure 1)
5	53°47'21.6"N, 20°27'54.0"E	Edge of the fresh pine forest <i>Peucedano-Pinetum</i> (potential vegetation: boreal spruce forest <i>Sphagno girgensohnii-Piceetum</i> ) and raised-bog community <i>Ledo-Sphagnetum magellanicum</i>
6	53°47'20.8"N, 20°27'56.9"E	Mixed old-growth spruce-birch forest (potential vegetation: boreal spruce forest <i>Sphagno girgensohnii-Piceetum</i> )
7	53°47'21.1"N, 20°27'51.2"E	Fresh pine forest <i>Peucedano-Pinetum</i> (potential vegetation: boreal spruce forest <i>Sphagno girgensohnii-Piceetum</i> )
<b>Redykajny</b>		
1	53°47'45.6"N, 20°27'10.7"E	Edge of the mixed forest (potential vegetation: continental mixed forest <i>Quercus roboris-Pinetum</i> ) and wet meadow
2	53°47'47.6"N, 20°27'11.4"E	Boggy pine forest <i>Vaccinio uliginosi-Pinetum</i> (Figure 2)
3	53°47'49.0"N, 20°27'10.0"E	Boggy pine forest <i>Vaccinio uliginosi-Pinetum</i>
4	53°47'50.8"N, 20°27'06.8"E	Alder swamp forest <i>Ribeso nigri-Alnetum</i>
5	53°47'44.1"N, 20°27'10.8"E	Group of old deciduous trees on the edge of a mixed forest (potential vegetation: continental mixed forest <i>Quercus roboris-Pinetum</i> ) and wet meadow
6	53°47'44.2"N, 20°27'17.1"E	Spruce forest (potential vegetation: continental mixed forest <i>Quercus roboris-Pinetum</i> )
7	53°47'42.7"N, 20°27'07.9"E	Multi-species group of trees on the edge of the reserve, by the mid-forest road
8	53°47'45.5"N, 20°27'05.7"E	Old alders on the edge of a mixed forest, next to a drainage ditch
9	53°47'46.9"N, 20°27'07.7"E	Fresh pine forest <i>Peucedano-Pinetum</i>
10	53°47'47.4"N, 20°27'07.0"E	Boggy pine forest <i>Vaccinio uliginosi-Pinetum</i>
11	53°47'47.9"N, 20°27'08.8"E	Edge of boggy pine forest <i>Vaccinio uliginosi-Pinetum</i> and raised-bog community <i>Ledo-Sphagnetum magellanicum</i>
12	53°47'47.5"N, 20°27'03.7"E	Mixed alder-birch forest (potential vegetation: swamp birch forest <i>Betuletum pubescentis</i> )
13	53°47'45.0"N, 20°27'02.2"E	Boreal spruce forest <i>Sphagno girgensohnii-Piceetum</i> (old-growth stand)
14	53°47'47.5"N, 20°26'58.2"E	Alder forest (potential vegetation: alder turf forest <i>Sphagno squarrosi-Alnetum</i> )
15	53°47'49.7"N, 20°26'56.4"E	Alder forest (potential vegetation: alder turf forest <i>Sphagno squarrosi-Alnetum</i> )
16	53°47'49.9"N, 20°26'52.9"E	Old elm tree on the edge of the reserve, near the forest path (potential vegetation: subboreal humid mixed forest <i>Quercus-Piceetum</i> )

ously mentioned knowledge and experience of the members of the research team), the number of species found in a given area positively correlates with the number of researchers and the amount of time spent in the field (Vondrák et al., 2016). Apart from the random (probabilistic) approach to data collection, which is particularly suitable for larger areas with a less diverse environment, in the practice of lichenological inventories of smaller areas, two methods of recording species can be distinguished. The first involves the relatively uniform penetration of the entire area (route or cartographic method; see Faliński, 1990), while in the second, the researchers focus their attention on places that are potentially important for the diversity of lichens, so-called biodiversity hotspots (Peterson & McCune, 2003; Vondrák et al., 2016, 2022). Both methods have their advantages and disadvantages, but

it would seem that in order to identify the largest number of species, in particular microlichens, the second approach is more efficient (Hofmeister et al., 2022; Vondrák et al., 2018). The optimal solution would be a combination of both methods (Ravera & Brunialti, 2013), but such a differentiated approach would hinder the analysis of the data obtained and the drawing of conclusions on their basis while also limiting the repeatability of the study. In assessing the results presented in this paper in terms of the applied research methodology, it should be noticed that a different method of collecting data was applied in the two different study periods. In the first period, the study was conducted using the route (topographic) method (see Faliński, 1990; Kubiak, 2008). It is difficult not to conclude that when using this method, the researcher has a tendency to avoid areas that

**Table 2** List of species recorded in the “Mszar” and “Redykajny” reserves in 1999–2001 and 2024. Numbers in the columns 4–5 correspond to the number of study sites in Table 1. Newly recorded species are in bold.

Species	1999–2001		2024		
	Mszar	Redykajny	Mszar	Redykajny	
	1	2	3	4	5
<i>Absconditella lignicola</i> Vězda & Pišút	–	+	5	3,12,13	
<i>Amandinea punctata</i> (Hoffm.) Coppins & Scheid.	+	–	–	1	
<i>Anaptychia ciliaris</i> (L.) Körb.	–	+	–	–	
<b><i>Anisomeridium polypori</i></b> (Ellis & Everh.) M.E. Barr	–	–	2	7,16	
<b><i>Bacidia arceutina</i></b> (Ach.) Arnold	–	–	1	7,13	
<i>Bacidia fraxinea</i> Lönnr.	–	+	–	7	
<i>Bacidia rubella</i> (Hoffm.) A. Massal.	–	+	–	7	
<i>Bacidina modesta</i> (Zwackh ex Vain.) S. Ekman	–	+	1–3	14	
<i>Biatora efflorescens</i> (Hedl.) Räsänen	–	+	2	14	
<b><i>Bryoria fuscescens</i></b> (Gyeln.) Brodo & D. Hawksw.	–	–	3	–	
<i>Buellia griseovirens</i> (Turner & Borrer ex Sm.) Almb.	+	+	2–5	3,4,7,10,14	
<b><i>Calicium trabinellum</i></b> (Ach.) Ach.	+	–	4	–	
<b><i>Candelariella efflorescens</i></b> R.C. Harris & W.R. Buck	–	–	4	4	
<i>Carbonicola anthracophila</i> (Nyl.) Bendiksby & Timdal	+	–	–	9	
<b><i>Catillaria nigroclavata</i></b> (Nyl.) J. Steiner	–	–	1	–	
<i>Cetraria sepincola</i> (Ehrh.) Ach.	+	–	–	–	
<b><i>Chaenotheca brachypoda</i></b> (Ach.) Tibell	–	–	–	7,13	
<b><i>Chaenotheca chlorella</i></b> (Ach.) Müll.Arg.	–	–	–	9	
<b><i>Chaenotheca chrysocephala</i></b> (Ach.) Th.Fr.	–	–	–	5,8,10	
<i>Chaenotheca ferruginea</i> (Turner ex Sm.) Mig.	+	+	1,2,4–7	4,5,7,9–11,14,15	
<i>Chaenotheca furfuracea</i> (L.) Tibell	–	+	–	4,13	
<i>Chaenotheca stemonea</i> (Ach.) Müll.Arg.	+	+	1,2,6,7	7,8	
<i>Chaenotheca trichialis</i> (Ach.) Th.Fr.	+	–	2,4,6,7	4,5,8,14,15	
<i>Chaenotheca xyloxena</i> Nádv.	–	+	4,7	9,10,13	
# <i>Chaenothecopsis pusiola</i> (Ach.) Vain.	–	+	–	9	
<i>Cladonia arbuscula</i> (Wallr.) Flot. em. Ruoss subsp. arbuscula	–	+	–	–	
<i>Cladonia cenotea</i> (Ach.) Schaer.	+	–	4,5,7	3,10	
<i>Cladonia chlorophaea</i> (Flörke ex Sommerf.) Spreng.	–	+	–	2,4,12	
<i>Cladonia coniocraea</i> (Flörke) Spreng.	+	+	1–7	1–5,7–15	
<i>Cladonia digitata</i> (L.) Hoffm.	+	+	2,4,5,7	1–5,9–11,14	
<i>Cladonia fimbriata</i> (L.) Fr.	+	+	2,4,5,7	1–3,7,9,10,12,13,15	
<i>Cladonia floerkeana</i> (Fr.) Flörke	+	–	7	10,11	
<i>Cladonia glauca</i> Flörke	+	–	–	9	
<i>Cladonia grayi</i> G. Merr. ex Sandst.	–	+	–	12	
<i>Cladonia macilenta</i> Hoffm.	+	+	1,4,5,7	2,3,5,10,15	
<b><i>Cliostomum leprosum</i></b> (Räsänen) Holien & Tønsberg	–	–	–	9	
<i>Coenogonium pineti</i> (Ach.) Lücking & Lumbsch	–	+	1–7	1–9,11–16	
<b><i>Diarthonis spadicea</i></b> (Leight.) Frisch, Ertz, Coppins & P.F. Cannon	–	–	2,4	4,5,7,8,13–16	
<i>Evernia prunastri</i> (L.) Ach.	+	+	1–3,5,7	5,6,11	
<i>Fuscidea arboricola</i> Coppins & Tønsberg	–	+	2,4	4,15	
<i>Fuscidea pusilla</i> Tønsberg	–	+	4,5	4,7,9,11,12,15	
<b><i>Graphis scripta</i></b> (L.) Ach.	–	–	1	8,14	
<i>Hypocenomyce scalaris</i> (Ach.) M. Choisy	+	+	1,4,5,7	2,3,5,7–9,11,14,15	

Continued on next page

Table 2 Continued.

Species	1999–2001		2024	
	Mszar	Redykajny	Mszar	Redykajny
<i>Hypogymnia physodes</i> (L.) Nyl.	+	+	1–5,7	2–7,9–15
<b><i>Hypogymnia tubulosa</i></b> (Schaer.) Hav.	–	–	2–4,7	6,9–11,13
<i>Imshaugia aleurites</i> (Ach.) S.L.F. Meyer	–	+	–	3,5,9,10
<i>Lecania cyrtella</i> (Ach.) Th.Fr.	–	+	–	11
<i>Lecania</i> cfr. <i>cyrtellina</i> (Nyl.) Sandst.	–	+	–	–
<b><i>Lecania naegelii</i></b> (Hepp) Diederich & van den Boom	–	–	2	–
<i>Lecanora argentata</i> (Ach.) Malme	–	+	–	–
<i>Lecanora carpinea</i> (L.) Vain.	–	+	–	–
<i>Lecanora chlarotera</i> Nyl.	–	+	–	7
<i>Lecanora conizaeoides</i> Nyl. ex Cromb.	+	+	1,3–5	3,5,10,11
<i>Lecanora expallens</i> Ach.	+	+	1,4,7	5,14,15
<i>Lecanora pulicaris</i> (Pers.) Ach.	+	+	1–4,7	3,4,6,9–12,14
<b><i>Lecanora sarcopidoides</i></b> (A. Massal.) Hedl.	–	–	4	–
<b><i>Lecanora stanislai</i></b> Guzow-Krzem., Łubek, Malíček & Kukwa	–	–	1,4	12,14,15
<i>Lecidea nylanderii</i> (Anzi) Th.Fr.	–	+	4	3,9–11
<i>Lecidella elaeochroma</i> (Ach.) M. Choisy	–	+	–	–
<b><i>Lecidella subviridis</i></b> Tønsberg	–	–	–	4,16
<b><i>Lepra albescens</i></b> (Huds.) Hafellner	–	–	–	14
<i>Lepra amara</i> (Ach.) Hafellner	–	+	–	5,8,14
<i>Lepraria eburnea</i> J.R. Laundon	–	+	–	4
<i>Lepraria elobata</i> Tønsberg	+	+	1–5,7	4,5,7,12–15
<i>Lepraria finkii</i> (B. de Lesd. ex Hue) R.C. Harris	–	+	–	5,7,8,13–16
<i>Lepraria incana</i> (L.) Ach.	+	+	1–7	1–5,7–9,11–16
<i>Lepraria jackii</i> Tønsberg	+	+	1–5,7	2–12,14
<b><i>Lepraria vouauxii</i></b> (Hue) R.C. Harris	–	–	–	4,7,12
<b><i>Lichenomphalia umbellifera</i></b> (L.) Redhead, Lutzoni, Moncalvo & Vilgalys	–	–	1	5,10,13
<i>Melanelixia glabratula</i> (Schaer.) O. Blanco et al.	+	+	1	7,14
<b><i>Melanelixia subaurifera</i></b> (Nyl.) O. Blanco et al.	–	–	2–4,7	7,10,11,14
<i>Melanohalea exasperatula</i> (Nyl.) O. Blanco et al.	–	+	7	5,11
<b><i>Micarea botryoides</i></b> (Nyl.) Coppins	–	–	1	–
<b><i>Micarea byssacea</i></b> (Th. Fr.) Czarnota, Guzow-Krzem. & Coppins	–	–	–	4,7
<i>Micarea denigrata</i> (Fr.) Hedl.	+	–	–	3,10,12
<i>Micarea melaena</i> (Nyl.) Hedl.	+	–	–	–
<b><i>Micarea micrococca</i></b> (Körb.) Coppins	–	–	1,4	9,11
<i>Micarea misella</i> (Nyl.) Hedl.	+	–	1,4,5	1,9,10,12
<i>Micarea</i> cfr. <i>nigella</i> Coppins	+	–	–	–
<i>Micarea nitschkeana</i> (J. Lahm ex Rabenh.) Harm.	–	+	–	–
<b><i>Micarea soralifera</i></b> Guzow-Krzemińska, Czarnota, Łubek & Kukwa	–	–	–	9,12
<b><i>Micarea viridileprosa</i></b> Coppins & van den Boom	–	–	7	12
<b>#<i>Microcalicium ahlneri</i></b> Tibell	–	–	–	9
<i>Nephromopsis chlorophylla</i> (Willd.) Divakar, Crespo & Lumbsch	–	+	2–4	–
<i>Ochrolechia bahusiensis</i> H. Magn.	–	+	2	4,7,8,12,14
<i>Ochrolechia microstictoides</i> Räsänen	–	+	–	–
<i>Palicella filamentosa</i> (Stirt.) Rodr. Flakus & Printzen	–	+	4	10,11,12

Continued on next page

Table 2 Continued.

Species	1999–2001		2024	
	Mszar	Redykajny	Mszar	Redykajny
<i>Parmelia submontana</i> Nád. ex Hale	–	+	–	–
<i>Parmelia sulcata</i> Taylor	+	+	1–5,7	1,3–7,9–11,14,15
<i>Parmeliopsis ambigua</i> (Wulfen) Nyl.	+	+	1,3,4,7	3,5,9,10,14
<b><i>Peltigera praetextata</i></b> (Flörke) Zopf	–	–	–	7
<i>Pertusaria coccodes</i> (Ach.) Nyl.	–	+	–	–
<i>Phaeophyscia orbicularis</i> (Neck.) Moberg	–	+	–	7
<i>Phlyctis argena</i> (Ach.) Flot.	–	+	2–4	4,7,12–15
<b><i>Physcia adscendens</i></b> (Fr.) H. Olivier	–	–	1,2	14
<i>Physcia stellaris</i> (L.) Nyl. subsp. <i>stellaris</i>	–	+	–	–
<i>Physcia tenella</i> (Scop.) DC.	–	+	1,2,4,7	2,4,10,11,14
<i>Physconia enteroxantha</i> (Nyl.) Poelt	–	+	–	–
<i>Placynthiella dasaea</i> (Stirt.) Tønsberg	+	+	1,2,4,5,7	2–5,10,12,13
<i>Placynthiella icmalea</i> (Ach.) Coppins & P. James	+	+	4,7	1,3–5,10,12
<i>Platismatia glauca</i> (L.) W.L. Culb. & C.F. Culb.	+	+	1–3	3,5,6,10,11,14
<b><i>Polycauliona polycarpa</i></b> (Hoffm.) Frödén, Arup & Söchting	–	–	–	14
<b><i>Porina aenea</i></b> (Wallr.) Zahlbr.	–	–	1,7	14
<i>Pseudevernia furfuracea</i> (L.) Zopf	+	+	1,3–5,7	3,9–11,14
<i>Pycnora sorophora</i> (Vain.) Hafellner	+	+	4,7	3,5,9–12
<i>Ramalina farinacea</i> (L.) Ach.	+	+	–	5,7
<i>Ropalospora viridis</i> (Tønsberg) Tønsberg	–	+	2,4	2,4,7,9,11–15
<i>Scoliciosporum chlorococcum</i> (Graeve ex Stenh.) Vězda	+	+	–	–
<i>Scoliciosporum sarothamni</i> (Vain.) Vězda	–	+	2,3	9,11
# <i>Stenocybe pullatula</i> (Ach.) Stein	–	+	–	4,14
<i>Thelidium minutulum</i> Körb.	–	+	–	–
<b><i>Toniniopsis separabilis</i></b> (Nyl.) Gerasimova & A. Beck	–	–	–	7
<i>Trapeliopsis flexuosa</i> (Fr.) Coppins & P. James	+	+	4,5,7	3,5,10
<b><i>Trapeliopsis gelatinosa</i></b> (Flörke) Coppins & P. James	–	–	4	–
<i>Trapeliopsis granulosa</i> (Hoffm.) Lumbsch	+	+	1,4,5,7	1,3,10,12,13
<i>Usnea hirta</i> (L.) Weber ex F.H. Wigg.	–	+	3	10,11
<i>Usnea subfloridana</i> Stirt.	+	+	4	–
<i>Vulpicida pinastri</i> (Scop.) J.-E. Mattson & M.J. Lai	+	–	4	3,10,12
<i>Violella fucata</i> (Stirt.) T. Sprib.	–	+	1,4,5,7	1–5,7,9–12,14
<i>Xanthoria parietina</i> (L.) Th.Fr.	–	+	1	–
<i>Xylopsora caradocensis</i> (Leight. ex Nyl.) Bendiksby & Timdal	–	+	–	–
<b>A total</b>	<b>42</b>	<b>76</b>	<b>71</b>	<b>93</b>
		<b>89</b>		<b>103</b>
			<b>121</b>	

# Non-lichenized fungi.

are difficult to access (as well as those that are permanently or seasonally inaccessible) or those that are assumed to be uninteresting in lichenological terms. This may lead to the omission of certain highly specialised species (Vondrák et al., 2024). In the case of studies conducted using the point method, this forces the researcher to conduct a systematic search of microhabitats and substrates, regardless of the preliminary assessment of their interest in lichenological terms. To minimize the subjectivity of the selection of research

sites, it is worthwhile to respect the principle of determining these in each of the identified types of habitats, including in their ecotonic zones, as was done during the second inventory in this study. It seems that the reason that many new species were identified, in particular those that belong to the so-called microlichens, at least in part, was due to the different procedures for data collection. Perhaps the number of their locations and the number of individuals increased during the analyzed period, which made their

identification easier in 2024. It seems that the appearance of some of the newly recorded species can be associated with an increase in the amount of deadwood among the spruce undergrowth. Quite characteristic communities of epiphytic lichens formed on the dead spruce branches composed both of lichens that had not been recorded previously (*H. tubulosa*, *B. fuscescens*, *M. subaurifera*) and species that were known from the previous inventory, but it would seem, had increased in numbers (*E. prunastri*, *Platismatia glauca*, *Pseudevernia furfuracea*, *U. hirta*). Numerous studies show that the population dynamics of organisms such as lichens are closely linked with the dynamics of forest substrates, either continuously linked or linked at certain stages of their individual development (Scheidegger & Werth, 2009). Such a role is played by deadwood in various forms, which for many epiphytes is a substitute substrate and a centre for dispersion of diaspores into the environment (Tanona & Czarnota, 2023). A separate category of dead wood is dead fallen tree trunks. Depending on the type of tree and local microclimatic conditions, this substrate may be colonized by specialized epixylites, which are different from those associated with deadwood in exposed places (especially in the case of dead, standing trees), which are present, for example, in the boggy pine forest community. As a result of the second inventory, several new species for the area were identified on the dead and fallen trunks of deciduous trees in shady and damp localities: *Ch. brachypoda*, *L. umbellifera*, *P. praetextata* and *Trapeliopsis gelatinosa*. The most significant risk for the forest lichen biota is the serious degradation or loss of habitat (Pykälä, 2019; Wolseley, 1995). In the case of the studied area, this type of threat was not observed, among other reasons, due to the habitat diversity of both reserves and their location within a large forest complex, which may act as a buffer against many potential threats. However, in both reserves, natural succession processes took place. Of particular importance are those that concern areas that are crucial for reasons of preservation – raised bogs and boggy pine forests. Despite the clear reduction in surface area of these communities, so far, significant changes in the species composition of lichens that are associated with them (mainly epiphytes of pine and epixylites associated with dead pine wood) have not been observed. It is worth stressing that the most recent study confirmed the presence in these communities of *C. trabinellum*, a species that is considered endangered (EN) in Poland. The species is not found in Olsztyn outside of the “Mszar” reserve (Kubiak, 2005) and is very rare in northern Poland (Cieśliński, 2003; Szczepańska et al., 2023). The assessment of environmental processes on the basis of individual species, especially ones that do not occur in large numbers, is of doubtful value. Its validity, however, increases in the case of a larger number of species, especially if these form a group with similar ecological requirements and functions performed in the ecosystem – the so-called functional traits (Ellis et al., 2021). As a result of the second inventory, in both reserves, a group of new species was identified, including epiphytic crustose lichens, which are considered characteristic of shady and damp deciduous forests (*Anisomeridium polypori*, *Diarthonia spadicea*, *G. scripta*, *Porina aenea*). These species undoubtedly came from the forest complex that surrounds both reserves, a complex that is dominated by stands of pine growing in relatively fertile habitats typical of oak-hornbeam forests (BDoL, 2024). This phenomenon can be interpreted as a

subtle signal of changes in the structure of conifer stands, primarily those growing at the edges of both peatland basins, which is expressed by an increasing share of deciduous species (including beech). Interestingly, the species are linked by the presence of the same photobiont from the genus *Trentepohlia* Mart. Aptroot and van Herk (2007), based on studies conducted in the Netherlands, showed that lichens containing this type of photobiont are increasing their range and numbers as a result of the global warming climate. Although comparative studies from the periods 1987–1989 and 2014–2015 conducted in the Białowieża Forest (one of the best-preserved forest complexes in the European lowlands) did not confirm a similar phenomenon (Łubek et al., 2021), it is much more likely in urbanized areas, which has so far been confirmed in studies of lichens in cities in western and southern Europe (Aptroot & van Herk, 2007; Munzi et al., 2014). As can be seen from the examples provided, the group of species recorded as new for the area in 2024 includes lichens with diverse morphology, habitat preferences, and reactions to environmental changes. It is worth emphasizing that threatened species are among them: *B. fuscescens*, *Ch. brachypoda*, *Ch. chlorella*, *G. scripta*, *H. tubulosa*, *L. sarcopidoides*, *L. umbellifera*, *P. praetextata*, *T. separabilis*, and *T. gelatinosa*.

### 3.3. Extinct and/or unconfirmed species

It should be noted that due to the previously described problems with complete identification of species, caution should be exercised when defining species as locally extinct. Undoubtedly, some species have disappeared from the research area, such as *C. arbuscula*, *A. ciliaris*, and *P. submontana*. The first species was found in the “Redykajny” reserve. A single, small thallus grew on a dead, strongly decomposed stump in a pine bog forest. Its extinction was probably caused by the natural succession of this substrate. The last two lichens were also recorded in the “Redykajny” reserve, where they grew on single, mature deciduous trees, *Acer platanoides* and *Populus tremula*. These trees had toppled, and as a result, the epiphytes inhabiting them died. These phorophytes are represented in minimal numbers in both reserves, creating a severe risk to the associated lichens. In the past, several familiar lichens were associated with the aforementioned phorophytes, which were not identified in 2024 (e.g., *Lecanora carpinea*, *Lecidella elaeochroma*, *Physcia stellaris*, *Phaeophyscia enteroxantha*). Among the rare and/or threatened lichens, the occurrence of only three species has not been confirmed: *C. sepincola*, *M. melaena*, and *P. coccodes*.

## 4. Conclusions

Knowledge of species and their distribution, especially in protected areas, is essential not only for scientific reasons but also for practical justification. It is crucial to undertake conservation measures and allow them to be adapted to the biology and ecology of individual species. However, it should be remembered that obtaining a complete list of lichen species in forests is very difficult, if not impossible. Therefore, it is important to use different data collection methods, allowing for obtaining a more complete list of species.

The conducted studies have shown that a relatively short period separating the two inventories is sufficient for changes



in relatively well-preserved forest communities that are significant enough to be reflected in the species composition of lichens. This confirms the generally known good bioindicative properties of lichens but, at the same time, forces the need for constant monitoring of the diversity of these organisms.

The different ecological requirements of lichens and their varying sensitivity to environmental changes can pose a significant challenge for nature conservation services and often require different compromises in their actions. It is important to establish conservation priorities, which should focus on selected species – rare and/or threatened, with well-known biology and ecology. Since these are usually stenoeccious species adapted to specific substrates and/or communities, the best solution is to protect their habitats.

## References

- Aptroot, A., Stapper, N. J., Košuthová, A., & van Herk, K. (2021). Lichens as an indicator of climate and global change. In T. M. Letcher (Ed.), *Climate change: Observed impacts on planet earth* (pp. 483–497). Elsevier. <https://doi.org/10.1016/B978-0-12-821575-3.00023-2>
- Aptroot, A., & van Herk, K. (2007). Further evidence of the effects of global warming on lichens, particularly those with *Trentepohlia* phycobionts. *Environmental Pollution*, 146, 293–298. <https://doi.org/10.1016/j.envpol.2006.03.018>
- BDoL. (2024). *Bank Danych o Lasach, Biuro Urządzania Lasu i Geodezji Leśnej Przedsiębiorstwo Państwowe* [Forest data bank, bureau for forest management and geodesy state enterprise]. <https://www.bdl.lasy.gov.pl/portal/>
- Bergamini, A., Scheidegger, C., Stofer, S., Carvalho, P., Davey, S., Dietrich, M., Dubs, F., Farkas, E., Groner, U., Kärkkäinen, K., Keller, C., Lokos, L., Lommi, S., Máguas, C., Mitchell, R., Pinho, P., Rico, V., Aragón, G., Robinson, A. M., & Watt, A. (2005). Performance of macrolichens and lichen genera as indicators of lichen species richness and composition. *Conservation Biology*, 19, 1051–1062. <https://doi.org/10.1111/j.1523-1739.2005.00192.x-i1>
- Boch, S., Prati, D., Hessenmöller, D., Schulze, E. D., & Fischer, M. (2013). Richness of lichen species, especially of threatened ones, is promoted by management methods furthering stand continuity. *Plos One*, 8, Article e55461. <https://doi.org/10.1371/journal.pone.0055461>
- Boggess, L. M., McCain, C. M., Manzitto-Tripp, E. A., Pearson, S. M., & Lendemer, J. C. (2024). Disturbance and diversity: Lichen species richness decreases with increasing anthropogenic disturbance. *Biological Conservation*, 293, Article 110598. <https://doi.org/10.1016/j.biocon.2024.110598>
- Chlebicki, A., Żarnowiec, J., Cieśliński, S., Klama, H., Bujakiewicz, A., & Załuski, T. (1996). Epixylites, lignicolous fungi and their links with different kinds of wood. In J. B. Faliński & W. Mułenko (Eds.), *Cryptogamous plants in the forest communities of Białowieża National Park (Project CRYPTO 3)*. *Phytocoenosis*, 8 (N.S.), *Archivum Geobotanicum*, 6, 75–110.
- Cieśliński, S. (2003). Atlas rozmieszczenia porostów (Lichenes) w Polsce Północno-Wschodniej [Atlas of the distribution of lichens (Lichenes) in North-Eastern Poland]. *Phytocoenosis* (N.S.), 15, *Supplementum Cartographiae Geobotanicae*, 15, 1–430.
- Cieśliński, S., Czyżewska, K., & Fabiszewski, J. (2006). Red list of extinct and threatened lichens in Poland. In Z. Mirek, K. Zarzycki, W. Wojewoda, & Z. Szelaż (Eds.), *Red list of plants and fungi in Poland* (pp. 71–89). W. Szafer Institute of Botany, Polish Academy of Sciences.
- Cieśliński, S., Czyżewska, K., Klama, H., & Żarnowiec, J. (1996). Epiphytes and epiphytism. In J. B. Faliński & W. Mułenko (Eds.), *Cryptogamous plants in the forest communities of Białowieża National Park (Project CRYPTO 3)*. *Phytocoenosis* (N.S.), 8, *Archivum Geobotanicum*, 6, 15–35.
- Cieśliński, S., & Fałtynowicz, W. (1993). Note from editors. In S. Cieśliński & W. Fałtynowicz (Eds.), *Atlas of the geographical distribution of lichens in Poland* (Vol. 1, pp. 7–8). W. Szafer Institute of Botany of Polish Academy of Sciences.
- Cislaghi, C., & Nimis, P. L. (1997). Lichens, air pollution and lung cancer. *Nature*, 387, 463–464. <https://doi.org/10.1038/387463a0>
- CRFOP. (2024). *Centralny Rejestr Form Ochrony Przyrody, Generalna Dyrekcja Ochrony Środowiska* [Central register of nature protection forms, General Directorate for Environmental Protection]. <https://crfop.gdos.gov.pl/CRFOP/index.jsf>
- Czarnota, P., & Guzow-Krzemińska, B. (2010). A phylogenetic study of the *Micarea prasina* group shows that *Micarea micrococca* includes three distinct lineages. *Lichenologist*, 42(1), 7–21. <https://doi.org/10.1017/S0024282909990211>
- Czerepko, J., Gawryś, R., Szymczyk, R., Pisarek, W., Janek, M., Haidt, A., Kowalewska, A., Piegdoń, A., Stebel, A., Kukwa, M., & Cacciatori, C. (2021). How sensitive are epiphytic and epixylic cryptogams as indicators of forest naturalness? Testing bryophyte and lichen predictive power in stands under different management regimes in the Białowieża forest. *Ecological Indicators*, 125, Article 107532. <https://doi.org/10.1016/j.ecolind.2021.107532>
- Czyżewska, K., & Cieśliński, S. (2003). Porosty – wskaźniki niżowych lasów puszczańskich w Polsce [Lichens – Indicators of lowland old-growth forests in Poland]. *Monographiae Botanicae*, 91, 223–239. <https://doi.org/10.5586/mb.2003.013>
- Ellis, C. J. (2012). Lichen epiphyte diversity: A species, community and trait-based review. *Perspectives in Plant Ecology, Evolution and Systematics*, 14(2), 131–152. <https://doi.org/10.1016/j.ppees.2011.10.001>
- Ellis, C. J., Asplund, J., Benesperi, R., Branquinho, C., di Nuzzo, L., Hurtado, P., Martínez, I., Matos, P., Nascimbene, J., Pinho, P., Prieto, M., Rocha, B., Rodríguez-Arribas, C., Thüs, H., & Giordani, P. (2021). Functional traits in lichen ecology: A review of challenge and opportunity. *Microorganisms*, 9(4), Article 766. <https://doi.org/10.3390/microorganisms9040766>
- Faliński, J. B. (1990). *Kartografia geobotaniczna. 2. Kartografia fitosocjologiczna* [Geobotanical cartography. 2. Phytosociological cartography]. Państwowe Przedsiębiorstwo Wydawnictw Kartograficznych.
- Faliński, J. B., & Mułenko, W. (Eds.). (1996). *Cryptogamous plants in the forest communities of Białowieża National*

- Park. Functional group analysis and general synthesis (Project CRYPTO 3). *Phytocoenosis*, 8 (N.S.), *Archivum Geobotanicum*, 6, 1–224.
- Fałtynowicz, W., Czarnota, P., Krzewicka, B., Wilk, K., Jabłońska, A., Oset, M., Ossowska, E. A., Śliwa, L., & Kukwa, M. (2024). *Lichens of Poland. A fifth annotated checklist*. W. Szafer Institute of Botany, Polish Academy of Sciences.
- Frati, L., & Brunialti, G. (2023). Recent trends and future challenges for lichen biomonitoring in forests. *Forests*, 14, Article 647. <https://doi.org/10.3390/f14030647>
- Gauslaa, Y. (2014). Rain, dew, and humid air as drivers of morphology, function and spatial distribution in epiphytic lichens. *Lichenologist*, 46(1), 1–16. <https://doi.org/10.1017/S0024282913000753>
- Gutowski, J. M., Bobiec, A., Ciach, M., Kujawa, A., Zub, K., & Pawlaczyk, P. (2023). *The afterlife of a tree*. WWF Polska.
- Guzow-Krzemińska, B., Czarnota, P., Łubek, A., & Kukwa, M. (2016). *Micarea soralifera* sp. nov., a new sorediate species in the *M. prasina* group. *Lichenologist*, 48(3), 161–169. <https://doi.org/10.1017/S0024282916000050>
- Guzow-Krzemińska, B., Maliček, J., Oset, M., Łubek, A., Tønsberg, T., & Kukwa, M. (2017). *Lecanora stanislai*, a new, sterile, usnic acid containing lichen species from Eurasia and North America. *Phytotaxa*, 329(3), 201–211. <https://doi.org/10.11646/phytotaxa.329.3.1>
- Hämäläinen, A., Fahrig, L., Strengbom, J., & Ranius, T. (2023). Effective management for deadwood-dependent lichen diversity requires landscape-scale habitat protection. *Journal of Applied Ecology*, 60(8), 1597–1606. <https://doi.org/10.1111/1365-2664.14429>
- Hofmeister, J., Vondrák, J., Ellis, C., Coppins, B., Sanderson, N., Maliček, J., Palice, Z., Acton, A., Svoboda, S., & Gloor, R. (2022). High and balanced contribution of regional biodiversity hotspots to epiphytic and epixylic lichen species diversity in Great Britain. *Biological Conservation*, 266, Article 109443. <https://doi.org/10.1016/j.biocon.2021.109443>
- Kubiak, D. (2005). Lichens and lichenicolous fungi of Olsztyn (NW Poland). *Acta Mycologica*, 40(2), 125–174. <https://doi.org/10.5586/am.2005.026>
- Kubiak, D. (2008). Porosty rezerwatów torfowiskowych „Mszar” i „Redykajny” na Pojezierzu Olsztyńskim [Lichens in peat bog reserves “Mszar” and “Redykajny” in Olsztyn Lake District]. *Parki Narodowe i Rezerваты Przyrody*, 27(1), 3–14.
- Loppi, S. (2014). Lichens as sentinels for air pollution at remote alpine areas (Italy). *Environmental Science and Pollution Research*, 21, 2563–2571. <https://doi.org/10.1007/s11356-013-2181-0>
- Łubek, A., Kukwa, M., Jaroszewicz, B., & Czortek, P. (2021). Shifts in lichen species and functional diversity in a primeval forest ecosystem as a response to environmental changes. *Forests*, 12(6), Article 686. <https://doi.org/10.3390/f12060686>
- Miller, J. E. D., Vilella, J., Stone, D., & Hardman, A. (2020). Using lichen communities as indicators of forest stand age and conservation value. *Forest Ecology and Management*, 475, Article 118436. <https://doi.org/10.1016/j.foreco.2020.118436>
- Munzi, S., Correia, O., Silva, P., Lopes, N., Freitas, C., Branquinho, C., & Pinho, P. (2014). Lichens as ecological indicators in urban areas: Beyond the effects of pollutants. *Journal of Applied Ecology*, 51(6), 1750–1757. <https://doi.org/10.1111/1365-2664.12304>
- Nash, T. H., & Gries, C. (1991). Lichens as indicators of air pollution. In O. Hutzinger (Ed.), *Air pollution. The handbook of environmental chemistry* (Vol. 4C, pp. 1–19). Springer. [https://doi.org/10.1007/978-3-540-47343-5\\_1](https://doi.org/10.1007/978-3-540-47343-5_1)
- Nimis, P. L., Scheidegger, C., & Wolseley, P. A. (Eds.). (2002). *Monitoring with lichens – Monitoring lichens* (pp. 1–4). NATO Science Series, Kluwer.
- Orange, A., James, P. W., & White, F. J. (2001). *Microchemical methods for the identification of lichens*. British Lichen Society.
- Peterken, G. F. (1996). *Natural Woodland – Ecology and conservation in Northern temperate regions*. Cambridge University Press.
- Peterson, E. B., & McCune, B. (2003). The importance of hotspots for lichen diversity in forests of Western Oregon. *Bryologist*, 106(2), 246–256. <http://www.jstor.org/stable/3244659>
- Phinney, N. H., Solhaug, K. A., & Gauslaa, Y. (2018). Rapid resurrection of chlorolichens in humid air: Specific thallus mass drives rehydration and reactivation kinetics. *Environmental and Experimental Botany*, 148, 184–191. <https://doi.org/10.1016/j.envexpbot.2018.01.009>
- Poelt, J., & Vězda, A. (1990). Über kurzlebige Flechten [On shortliving lichens]. *Bibliotheca Lichenologica*, 38, 377–394.
- Pykälä, J. (2019). Habitat loss and deterioration explain the disappearance of populations of threatened vascular plants, bryophytes and lichens in a hemiboreal landscape. *Global Ecology and Conservation*, 18, Article e00610. <https://doi.org/10.1016/j.gecco.2019.e00610>
- Ravera, S., & Brunialti, G. (2013). Epiphytic lichens of a poorly explored National Park: Is the probabilistic sampling effective to assess the occurrence of species of conservation concern? *Plant Biosystems*, 147, 115–124. <https://doi.org/10.1080/11263504.2012.736425>
- Scheidegger, C., & Stofer, S. (2015). Bedeutung alter Wälder für Flechten: Schlüsselstrukturen, Vernetzung, ökologische Kontinuität [The importance of old-growth forests for lichens: Keystone structures, connectivity, ecological continuity]. *Schweizerische Zeitschrift für Forstwesen*, 166(2), 75–82. <https://doi.org/10.3188/szf.2015.0075>
- Scheidegger, C., & Werth, S. (2009). Conservation strategies for lichens: Insights from population biology. *Fungal Biology Reviews*, 23(3), 55–66. <https://doi.org/10.1016/j.fbr.2009.10.003>
- Sillet, S. C., & Antoine, M. E. (2004). Lichens and bryophytes in forest canopies. In M. R. Lowman & H. B. Rinker (Eds.), *Forest canopies* (pp. 151–174). Elsevier Academic Press. <https://doi.org/10.1016/B978-012457553-0/50013-7>
- Solon, J., Borzyszkowski, J., Bidłasik, M., Richling, A., Badora, K., Balon, J., Brzezińska-Wójcik, T., Chabudziński, Ł., Dobrowolski, R., Grzegorzczak, I., Jodłowski, M., Kistowski, M., Kot, R., Krąż, P., Lechnio, J., Macias, A., Majchrowska, A., Malinowska, E., Migoń, P., ... Ziąja, W. (2018). Physico-geographical mesoregions of Poland: Verification and adjustment of boundaries on the basis of contemporary spatial data. *Geographia Polonica*, 91(2), 143–170. <https://doi.org/10.7163/GPol.0115>

- Stanton, D. E., Ormond, A., Koch, N. M., & Colesie, C. (2023). Lichen ecophysiology in a changing climate. *American Journal of Botany*, *110*(2), Article e16131. <https://doi.org/10.1002/ajb2.16131>
- Szczepańska, K., Kubiak, D., Ossowska, E. A., Kukwa, M., Jaskólska, J., Kowalewska, A., Schiefelbein, U., Bohdan, A., Kepel, A., Sęktas, M., Szymczyk, R., Hachułka, M., Rutkowski, K., Smoczyk, M., Zalewska, A., Piegdoń, A., & Romanow-Pękał, E. (2023). Materiały do rozmieszczenia porostów i grzybów naporostowych Polski, 3 [Materials for the distribution of lichens and lichenicolous fungi in Poland, 3]. *Wiadomości Botaniczne*, *67*. <https://doi.org/10.5586/wb.671>
- Tanona, M., & Czarnota, P. (2023). The response of lichens inhabiting exposed wood of spruce logs to post-hurricane disturbances in Western Carpathian forests. *Fungal Ecology*, *63*, Article 101228. <https://doi.org/10.1016/j.funeco.2023.101228>
- Tripp, E. A., Lendemer, J. C., & McCain, C. M. (2019). Habitat quality and disturbance drive lichen species richness in a temperate biodiversity hotspot. *Oecologia*, *190*, 445–457. <https://doi.org/10.1007/s00442-019-04413-0>
- Verey, M. (2017). Theoretical analysis and practical consequences of adopting a model ATPOL grid as a conical projection defining the conversion of plane coordinates to the WGS 84 ellipsoid. *Fragmenta Floristica et Geobotanica Polonica*, *24*(2), 469–488.
- Vondrák, J., Hofmeister, J., Ellis, C., Coppins, B., Sanderson, N., Malíček, J., Palice, Z., Acton, A., & Svoboda, S. (2024). Bringing rare species into the open: They significantly contribute to species richness in British lichen biodiversity hot-spots. *Authorea*, *30*. <https://doi.org/10.22541/au.170665053.30701190/v1>
- Vondrák, J., Malíček, J., Palice, Z., Bouda, F., Berger, F., Sanderson, N., Acton, A., Pouska, V., & Kish, R. (2018). Exploiting hot-spots; effective determination of lichen diversity in a Carpathian virgin forest. *Plos One*, *13*(9), Article e0203540. <https://doi.org/10.1371/journal.pone.0203540>
- Vondrák, J., Malíček, J., Palice, Z., Coppins, B., Kukwa, M., Czarnota, P., Sanderson, N., & Acton, A. (2016). Methods for obtaining more complete species lists in surveys of lichen biodiversity. *Nordic Journal of Botany*, *34*(5), 619–626. <https://doi.org/10.1111/njb.01053>
- Vondrák, J., Svoboda, S., Malíček, J., Palice, Z., Kocourková, J., Knudsen, K., Mayrhofer, H., Thüs, H., Schultz, M., Košnar, J., & Hofmeister, J. (2022). From Cinderella to Princess: An exceptional hotspot of lichen diversity in a long-inhabited central-European landscape. *Preslia*, *94*, 143–181. <https://doi.org/10.23855/preslia.2022.143>
- von Hirschheydt, G., Kéry, M., Ekman, S., Stofer, S., Dietrich, M., Keller, C., & Scheidegger, C. (2024). Occupancy model reveals limited detectability of lichens in a standardised large-scale monitoring. *Journal of Vegetation Science*, *35*, Article e13255. <https://doi.org/10.1111/jvs.13255>
- Wolseley, P. A. (1995). A global perspective on the status of lichens and their conservation. *Berichte der Eidgenössischen Forschungsanstalt für Wald, Schnee und Landschaft*, *70*(1), 11–27.
- Wroblewski, A., Ernst, S., Weber, T., & Delach, A. (2023). The impact of climate change on endangered plants and lichen. *PLOS Climate*, *2*(7), Article e0000225. <https://doi.org/10.1371/journal.pclm.0000225>