

Article ID: 196668
DOI: [10.5586/asbp/196668](https://doi.org/10.5586/asbp/196668)

Publication History

Received: 2024-09-27
Accepted: 2024-11-30
Published: 2025-05-27

Handling Editor

Anna Mikula; Polish Academy of Sciences Botanical Garden – Center for Biological Diversity Conservation in Powsin, Warsaw, Poland; <https://orcid.org/0000-0002-6846-3090>

Authors' Contributions

EW-C: Research concept and design; EW-C, AS-R: Collection and/or assembly of data; EW-C, AS-R: Data analysis and interpretation; EW-C, AS-R: Writing the article; EW-C, AS-R: Critical revision of the article; EW-C, AS-R: Final approval of the article

Funding

This research was financially supported through a statutory funds from the Polish Ministry of Education and Science.

Competing Interests

No competing interests have been declared.

Copyright Notice

© The Author(s) 2025. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits redistribution, commercial and noncommercial, provided that the article is properly cited.

RESEARCH PAPER

Plasmodial tapetum, orbicules, and ultrastructure of *Myosotis scorpioides* L. (Boraginaceae) pollen grains

Elżbieta Weryszko-Chmielewska , Aneta Sulborska-Różycka 

Department of Botany and Plant Physiology, University of Life Sciences in Lublin, Akademicka 15, 20-950 Lublin, Poland

* Corresponding author. Email: aneta.sulborska@up.lublin.pl

Abstract

Myosotis scorpioides L. produces one of the smallest pollen grains. To date, the morphology and structure of the sporoderm of these grains have been studied, but the ultrastructure of their protoplasts has not been sufficiently investigated. The aim of the study was to analyze the structure of the anther wall, determine the tapetum type, and examine the size and subcellular aspects of *M. scorpioides* pollen grains. The study was carried out with the use of light, scanning, and transmission electron microscopy techniques. In the pollen maturity phase, the epidermis, the endothecium with reinforced walls, and the remains of the middle layer and cell walls of the plasmodial tapetum were distinguished in the anther wall. The characteristic traits of *M. scorpioides* anthers include the plasmodial tapetum and the presence of polygonal orbicules. Plasmodial tapetum protoplasts were observed in close proximity to the pollen grains. The orbicules, fused with the sporoderm, were present in both the tapetum cell remains and on the surface of the pollen grains. The length of the polar and equatorial axes of the grains was 7.5 µm and 3.2 µm, respectively. The subcellular characteristics of the pollen grains included a vegetative cell with a lobulated nucleus, numerous plastids surrounded by a rough endoplasmic reticulum, numerous mitochondria, rough endoplasmic reticulum profiles, various-sized vesicles, highly numerous ribosomes, and two sperm cells with a small amount of cytoplasm. The plastids and mitochondria were spherical and substantially smaller than those present in somatic cells.

Keywords

forget-me-not; anthers; loculus; periplasmodium; sporomorphs; protoplasts

1. Introduction

The genus *Myosotis* L. (Boraginaceae) comprises approximately 100 species growing in two main different centers: Eurasia and New Zealand (Chacón et al., 2016; Luebert et al., 2016). *Myosotis scorpioides* L. grows in the wild across Europe and in some areas of Asia (Siberia, Mongolia, the Caucasus) (Germplasm Resources Information Network [GRIN]) In Poland, the species is found across the country (https://www.atlas-roslin.pl/pelna/gatunki/Myosotis_palustris.htm). *M. scorpioides* plants mostly occur on water banks, wet meadows, thickets, bog springs, and peat bogs (Kłosowski & Kłosowski, 2006; Podbielkowski & Sudnik-Wójcikowska, 2003), where they form large knee-high colonies (Fletcher & Tomblin, 2005). The species prefers moderately sunny locations and wet humus- and nutrient-rich soils (https://www.atlas-roslin.pl/pelna/gatunki/Myosotis_palustris.htm). It is also grown as an ornamental plant (Podbielkowski & Sudnik-Wójcikowska, 2003).

The genus *Myosotis* belongs to the subfamily *Cynoglossoideae* and tribe *Myosotideae* Rchb. (Weigend et al., 2016). *M. scorpioides* produces light blue flowers with flat-spread corolla lobes. Its buds and young flowers are pink (Podhajská & Rivola, 1992). The plant blooms from May to September. The flowers are entomophilous or self-pollinating (Rutkowski, 2006).

Myosotis pollen is very small and represents the smallest pollen grains known (Stanley & Linskens, 1974). Most *Cynoglossoideae*, including *Myosotis* plants, produce heterocolpate pollen characterized by the presence of two alternately arranged types of apertures: true apertures (colpori) and pseudoapertures (pseudocolpi) (Hargrove & Simpson, 2003; Weigend et al., 2016). It has been shown that heterocolpate pollen grains are not found in the other Boraginaceae subfamilies (Díez & Valdés, 1991).

The morphology of pollen grains in the genus *Myosotis* is highly diverse (Beug, 2004; Meudt, 2016). Large differences

have been found in their size and shape and in the number of apertures (Díez & Valdés, 1991; Noroozi et al., 2022).

The morphology and ultrastructure of the sporoderm of *Myosotis palustris* pollen grains have been thoroughly investigated by Volkova et al. (2013). The development and structure of pollen grains in *M. azorica* and *M. laxa* growing in Argentina were presented by Strittmatter and Galati (2001). Meudt (2016) carried out comparative studies of the morphology of pollen grains in 30 different taxa of *Myosotis* growing in New Zealand. The development of the sporoderm in *M. scorpioides* was studied in Russia by Volkova et al. (2017). There is no sufficient data in the literature regarding the ultrastructure of the cytoplasm in *Myosotis* pollen grains.

The aim of the study was to analyze the structure of the anther wall, identify the tapetum type, and examine the size and subcellular aspects of *M. scorpioides* pollen grains, as there are no data on this issue in the available literature.

2. Materials and methods

2.1. Plant material

The samples were collected from *Myosotis scorpioides* L. plants growing in their natural localities in Krężnica Jara (51°09'04" N, 22°28'52" E) in 2022–2023. The botanical identification of the specimens was carried out by taxonomy specialist Professor Bożena Denisow. The plant specimens were also compared with samples (voucher specimen no. 152) deposited in the herbarium of the Department of Botany and Plant Physiology, University of Life Sciences in Lublin (Poland).

2.2. Pollen grain size – Light microscopy (LM) studies

Glycerine jelly preparations of pollen grains were made, and the length of their polar (P) and equatorial (E) axes was measured ($n = 50$) with the use of a Nikon Eclipse 400 light microscope (Nikon, Tokyo, Japan).

2.3. Morphology of pollen grains – Scanning electron microscopy (SEM) studies

To examine the pollen morphology using a scanning electron microscope (SEM), pollen grains were collected from the flowers on flowering days 1–2. The methodology was based on a simplified microscopic preparatory approach. The pollen grains were placed on the surface of stubs covered with carbon tape and sputtered with gold using an Emitech SC 7640 sputter coater (Polaron, Newhaven, East Sussex, the UK). The observations of the shape and surface of the pollen grains and the documentation were made using a TESCAN/VEGA LMU (Tescan, Brno, the Czech Republic) scanning electron microscope (an accelerating voltage of 30 kV) equipped with a TESCAN attachment for digital processing of microscopic images.

2.4. Ultrastructure of pollen grains – Transmission electron microscope (TEM) studies

The pollen grains were fixed in 4% glutaraldehyde in phosphate buffer (pH 7.2; 0.1 M) at 4 °C for 12 h, washed three times in phosphate buffer, and dehydrated in an ethanol

series. The samples were post-fixed in a 1% osmium tetroxide solution at 0 °C for 1.5 h and washed three times in distilled water. In the next step, they were dehydrated in a graded ethanol series and embedded in LR white resin (LR white acrylic resin, medium grade, Sigma-Aldrich). Following polymerization at 60 °C, ultrathin sections (60–70 nm) of the embedded material were obtained for TEM examinations (copper grids with a 300 square mesh were used) using a Reichert Ultracut S ultramicrotome and a glass knife. Finally, the pollen grains were stained with 0.5% uranyl acetate and post-stained with 0.5% lead citrate (Reynolds, 1963). The ultrastructure of the pollen grains was analyzed with the use of a JEM 1400 (JEOL Cp., Japan) transmission electron microscope at an accelerating voltage of 120 kV. Photographic documentation was made using an 11 Magapixel TEM Camera MORADA G2 (EMSIS GmbH, Münster, Germany).

The diameter of plastids and mitochondria ($n = 20$ each) and the height of orbicules ($n = 20$) were measured (5 samples).

3. Results

3.1. Anther structure

M. scorpioides produces actinomorphic gamopetalous flowers. Their equally sized stamens (5) are fused with the upper part of the corolla tube (Figure 1A). The length of the anther is 0.7–0.9 mm ($0.8 \mu\text{m} \pm 0.08$; mean $\pm$ SD), and its diameter at the widest point is 0.3–0.4 mm ($0.37 \mu\text{m} \pm 0.04$; mean $\pm$ SD). The apical part of the anther ends with a slightly centrifugally bent appendix (Figure 1A–B). The anthers are latrorse and dehisce by means of a longitudinal slit (Figure 1A). The pollen grains are much smaller than the anther epidermis cells (Figure 1C–D). The surface of the epidermis cells is covered by a cuticle with characteristic ornamentation (Figure 1D).

The cross-section of the anther wall shows three layers: a one-layered epidermis covered by cuticle, the endothecium with large cells with bar thickenings, and the remains of the middle layer and tapetum tangential walls (Figure 2A). The size of the epidermis cells in the stomium region is substantially smaller than the other cells of this tissue (Figure 2B). In the outer wall of the epidermis cells, there are electron-translucent and electron-dense zones with numerous small droplets (Figure 2C), whose presence may indicate the secretory properties of this tissue.

The inner surface of the anther wall is covered by orbicules (Figure 2A–B). They are also present on the walls of the endothecium radial walls (Figure 2B) and on the surface of cells forming a septum between the loculi. It seems that the orbicules are associated with the tapetum cell remains present on the inner locule surface (as seen in Figure 3A–B) (Figure 2D). The orbicules are polygon-like in cross-section with an electron-opaque cavity and an electron-translucent orbicule wall (Figure 3A–B). Their size is in the range of $0.42\text{--}0.67 \mu\text{m}$ ($0.55 \mu\text{m} \pm 0.07$ on average). In some areas, the orbicules and the pollen grain sporoderm are fused (Figure 3B). Plasmodial tapetum protoplasts are visible from the microspore stage to the 3-celled microgametophyte stage; they surround the pollen grains, forming a periplasmodium layer (Figure 3C–E).




Figure 1 Fragments of *Myosotis scorpioides* L. flowers and stamens (SEM). Upper part of the flower with visible fornicies and stamens fused with the corolla tube (A); anther in a lateral position; centrifugally bent connective process in the apical part (yellow asterisk) (B); fragment of a mature anther after the release of pollen grains (red arrows) with a visible loculus (C); anther surface with visible cuticle ornamentation on the outer wall of epidermis cells and mature pollen grains (D). c – corolla tube; f – fornicies; lo – loculus; s – stamens. Scale bars: (A) = 200 μm , (B) = 100 μm , (C) = 20 μm , (D) = 5 μm .

3.2. Pollen grain morphology

M. scorpioides pollen grains are very small and dumbbell-shaped with a constriction in the equatorial plane (Figure 3F). The average length of the polar axis (P) is 7.5 μm (± 0.16), the length of the equatorial diameter (E) is 3.2 μm (± 0.20), and the diameter of the pollen grain at the widest point of the pole is 4.5 μm (± 0.15). The grains are heterocolpate with three pseudocolpi and three alternately arranged colpi (Figure 3F). The pores are fused with colpi and located in the equatorial plane. Given the number and type of apertures, *M. scorpioides* pollen grains are classified as tricolporate. The structure of their poles allows the classification of the grains as isopolar and subhexagonal in the polar view. At each pole, there is a pseudoaperture with granular ornamentation (Figure 3G). The exine ornamentation in the intercolpi is psilate, while the margins of the colpi are granular (Figure 3H).

3.3. Ultrastructure of pollen grains

The ectexine and endexine in the intercolpi are very thin, whereas the endexine near the endoaperture is thick (Figure 4A). The intine is thick only on the endoaperture. Pollenkit is visible on the exine surface and among the bacula (Figure 4B). Mature pollen grains are 3-celled (Figure 4A–C). Two sperm cells are enclosed in the vegetative cell. The sperm cell has a very small amount of cytoplasm. The nuclei of the sperm cells are slightly lobulated and unilaterally flattened (Figure 4C–D). Many organelles are situated in the cytoplasm of the vegetative cell, which has a highly lobulated nucleus (Figure 4C–E). In the polar regions, there are plastids surrounded by rough endoplasmic reticulum (RER) profiles (Figure 5A–C). The size of the plastids is in the range of 0.38–0.50 μm (0.43 $\mu\text{m} \pm 0.05$ on average). The plastids contain tubule-like thylakoids (Figure 5A–C). Very numerous ribosomes and RER profiles are present in the entire vegetative cell (Figure 4C–D, Figure 5C). Mitochondria, circular in cross-




Figure 2 Ultrastructure of *M. scorpioides* L. anther wall (TEM). Cross-section of an anther wall with a visible epidermis layer, endothecium with fibrous bands, remains of middle layer (red asterisks), and orbicules (blue arrows) on the surface of tapetum tangential walls (A); anther wall cells located adjacent to the stomium: epidermis, small stomium epidermis cell, endothecium, orbicules (blue arrows), and crushed cells during stomium rupture (B); fragment of an epidermal cell with cell wall details: cuticle, cell wall with zones of electron-dense, electron-transparent, and small droplets (white arrows) (C); deformed cells forming the septum between the loculi after anther rupture; blue arrows indicate orbicules (D). cc – crushed cells; cu – cuticle; cw – cell wall; en – endothecium; ep – epidermis; sc – stomium cells. Scale bars: (A–B) = 5 μ m, (D) = 2 μ m, (C) = 1 μ m.

section, are concentrated in the equatorial region of the pollen grains (Figure 4A, Figure 4D–E). Their diameter is in the range of 0.19–0.20 μ m (0.19 μ m $\pm$ 0.01 on average). Numerous different-sized vesicles are visible in the cytoplasm (Figure 4D, Figure 5A). The pollen grains germinate through colpi (Figure 5D).

4. Discussion

The current study presents details of the structure of the mature anther wall in *M. scorpioides*, including details of the stomium and identification of the tapetum type. At this stage of development, the anther locus was surrounded only by the epidermis and the endothecium, as well as the remains of the middle layer and tapetum cell walls, with orbicules visible on their surface. The parietal layers between the endothecium and the tapetum were crushed and collapsed. Similar changes in mature anthers were highlighted by Esau (1973). Secondary thickening of endothecium cell walls was visible as ledges both on the internal tangential walls and on the radial walls. Garcia (2002) distinguished four main types of endothelial secondary thickening: (1) annual rib type, (2) helical rib type, (3) reticulate ribs, and (4) palmate ribs. The present observations suggest that the thickening in the *M. scorpioides* endothecium can be classified as the annual rib type.

We observed that the stomium region of the dehiscient *M. scorpioides* anthers after pollen release exhibited the presence of

well-preserved small epidermis cells, which are typical of this part of the anther. Close to these cells, there were crushed and deformed septum cells that previously separated the two loculi. As reported by Wilson et al. (2011), the septum in the mature anther is enzymatically lysed and undergoes a programmed cell death-like breakdown. In turn, the stomium rupture is a consequence of stresses associated with pollen swelling and anther dehydration. Ligno-cellulosic secondary thickening in endothecium walls, which is well formed in *M. scorpioides* anthers, also plays an important role in the opening of anthers.

The present study shows that *M. scorpioides* has a plasmodial tapetum, characterized by the release of tapetum cell protoplasts from cell walls and their penetration into the locus, where they surround and nourish the forming pollen grains. The protoplasts fuse to form a multinucleate periplasmodium (Furness, 2008; Pacini, 1996). During the development of pollen grains, plasmodia gradually disappear (Esau, 1973). Other authors have reported the presence of a secretory tapetum in most Boraginaceae representatives (Izmailow & Biskup, 2003; Johri et al., 1992; Rao & Rao, 1992). The secretory tapetum is predominant in angiosperms (Furness & Rudall, 2001; Pacini et al., 1985). In this tapetum type, a layer of tapetal cells surrounds the anther locus. In previous studies, the tapetum in *Myosotis azorica*, *M. laxa* (Strittmatter & Galati, 2001), and *M. scorpioides* (Volkova et al., 2017) was classified as the secretory type. A secretory tapetum was also




Figure 3 Inner part of the *M. scorpioides* L. anther wall surrounding the loculus, pollen grains, and plasmodial tapetum in the 3-celled pollen grain phase (TEM) (A–D). Heterocolpate pollen grains of *M. scorpioides* L. viewed in the scanning electron microscope (SEM) (F–H). Remains of tapetum cell walls and deformed cells of the intermediate layer and orbicules (blue arrows) (A); orbicules fused with the pollen grain wall (black arrows) (B); plasmodial tapetum protoplasts located in close proximity to pollen grains (C); pollen grains surrounded by the periplasmodium (D–E); Pollen grains with equatorial constrictions and furrows with various lengths (F); pseudoporus (yellow arrow) with granular exine ornamentation at the poles of the pollen grains (G); pollen grain with a longer pseudocolpus and a short colpus containing a porus (purple arrow); granular exine ornamentation on the edges of the furrows (H). pg – pollen grains; pt – plasmodial tapetum; pp – periplasmodium; ps – pseudocolpus; sc – short colpus. Scale bars: (D, G) = 2 μ m, (C, E, H) = 1 μ m, (A, B) = 0.5 μ m.

identified in *Symphytum orientale* (Boraginaceae) (Vardar & Yavuz, 2018), whereas two types of tapetum, i.e. secretory and plasmodial, were present in *Symphytum officinale* (Gabarayeva et al., 2011). It seems that *M. scorpioides* may be characterized by the occurrence of different types of tapetum, i.e., secretory and plasmodial.

In the present study, the tapetum penetrated the anther loculus at the microspore stage and remained there until the later stage of pollen development. Similar observations of the functioning of the plasmodial tapetum in *Butomus* (Butomaceae) were reported by Fernando and Cass (1994). Plasmodial tapetum has also been described in many other




Figure 4 Ultrastructure of *M. scorpioides* L. pollen grains (TEM). Longitudinal section of a pollen grain (A); fragment of a pollen grain with visible sporoderm and pollenkitt (green arrow) (B); submedial parts of protoplasts with sperm cells, plastids, numerous mitochondria, vesicles, RER profiles, and very numerous ribosomes (arrowheads) (C–D); lobulated vegetative nucleus with a nucleolus, sperm cells, numerous mitochondria, and plastids (E). mc – sperm cells; m – mitochondria; nu – nucleolus; pl – plastids; rer – rough endoplasmic reticulum; ve – vesicles; vn – vegetative nucleus. Scale bars: (A) = 1 μ m, (B–E) = 0.5 μ m.

Eudicots. In Asteraceae (e.g. *Helianthus annuus*) (Çetinbaş & Ünal, 2015) and Caprifoliaceae, plasmodial tapeta predominate (Furness, 2008; Pacini, 1996). This tapetum type has also been found in, e.g., Apiaceae, Berberidaceae, Gentianaceae, Oleaceae, Verbenaceae (Furness, 2008), and Alismataceae (*Hydrocleys nymphoides*, *Alisma plantago-aquatica*, and *Sagittaria montevidensis*) (Nicolau et al., 2024).

In the *M. scorpioides* anthers, polygonal-shaped orbicules were observed on the surface of the tapetum cell remains and on the surface of the pollen grains. This morphological

type of orbicules occurs in a few plant species, e.g., in some Piperales (Oak et al., 2022). Most often, orbicules have a spherical or nearly spherical shape (Ruggiero & Bedini, 2020; Vinckier & Smets, 2002a, 2002b). To date, great diversity in the shape, size, density, and ornamentation of orbicules has been observed in different families (Galati, 2003; Oak et al., 2022; Ruggiero & Bedini, 2018, 2020; Verstraete et al., 2014). Orbicules are sporopollenin particles involved in the formation of the sporoderm of pollen grains (Huysmans et al., 1998). The fusion of the orbicules with the cell wall of pollen




Figure 5 Fragments of the longitudinal section (a, b) and cross-section (c) of *M. scorpioides* L. pollen grains and a pollen grain with a germinating pollen tube (d) (TEM). Numerous circular plastids with thylakoids surrounded by RER profiles, vesicles, RER profiles, very numerous ribosomes (arrowheads) (A–C); pollen tube germinating from the equatorial part of the pollen grain (D). pl – plastids; pt – pollen tube; rer – rough endoplasmic reticulum; ve – vesicles. Scale bars: (D) = 1 µm, (A–C) = 0.5 µm.

grains in the anthers of *M. scorpioides*, which was shown in the present study by transmission electron microscopy, may confirm this information.

As shown by literature data, orbicules occur mainly in anthers that contain a secretory tapetum (Huysmans et al., 1998; Verstraete et al., 2014; Vinckier & Smets, 2003). Therefore, it may be assumed that the occurrence of orbicules in *M. scorpioides* anther loculi where the plasmodial tapetum was present is an exception. The unique species in this respect reported previously included *Gentiana acaulis* L. (Lombardo & Carraro, 1976; Vinckier & Smets, 2002a; 2002b) and *Tradescantia virginiana* (Tiwari & Gunning, 1986), in which orbicules were accompanied by a plasmodial tapetum as well. Galati et al. (2007) reported the presence of orbicules in the anthers of *Modiolastrum malvifolium* (Malvaceae), which had an invasive non-syncytial tapetum classified as intermediate between secretory and plasmodial tapetum. The authors found that the presence of orbicules is not associated with the tapetum type, as they occur in both the secretory and plasmodial tapetum.

The shape of the polygonal orbicules present in the *M. scorpioides* studied differed from the shape of orbicules found in other *Myosotis* species. Strittmatter and Galati (2001)

reported the presence of spherical and subspherical orbicules in *M. azorica* and *M. laxa*. These data indicate diversity in the morphology of orbicules in different species of the genus.

The size of the orbicules in *M. scorpioides* was in the range of 0.42–0.67 µm, i.e. they can be classified as small in comparison with other literature data. Vinckier and Smets (2003) reported the size of orbicules in representatives of the family Gentianaceae in the range of 0.14–2.98 µm. As shown by Ruggiero and Bedini (2020), the size of orbicules in 34 species from different families ranged from 0.06 to 2.56 µm. In turn, Oak et al. (2022) reported a range of 0.24–2.79 µm in 12 Piperales species. As reported by Ruggiero and Bedini (2020), there is an evolutionary trend towards a reduction in the size of orbicules in angiosperms. Tiwari and Gunning (1986) found that orbicules in anthers with ameboid tapeta are smaller than in secretory tapeta.

The present study indicated the presence of three cells in mature pollen grains in *M. scorpioides*: a vegetative cell and two sperm cells located therein. Similarly, 3-celled pollen grains were described in *M. azorica* and *M. laxa* by Strittmatter and Galati (2001). In their study on the development of pollen grains in *M. scorpioides*, Volkova et al. (2017) carried

out a detailed analysis of the development of the sporoderm in this species, but without information on the internal structure of pollen grains.

The analyses of the ultrastructure of *M. scorpioides* pollen grains revealed the presence of slightly lobulated nuclei and small amounts of cytoplasm in the sperm cells. In turn, the nucleus of the vegetative cell was intensively lobulated. The cytoplasm of the vegetative cell contained numerous plastids surrounded by the RER. The plastids had a system of internal membranes. There were also mitochondria, RER profiles, very numerous ribosomes, and different-sized vesicles. Some of these structures, i.e., proplastids, RER profiles, and mitochondria, were observed in the vegetative cells of *M. azorica* and *M. laxa* (Strittmatter & Galati, 2001). Noteworthy are the exceptionally small sizes of plastids (0.38–0.50 µm) in the *M. scorpioides* pollen grains. Fujiwara et al. (2010) described plastids in pollen grains and growing tubes from transgenic *Arabidopsis thaliana* reaching a length of 0.3–3.1 µm (1.4 ± 0.4 µm on average). In turn, plastids with a diameter of 4–6 µm were found in somatic cells of angiosperms (Evert, 2006). Boussardon et al. (2023) described plastids in mesophyll cells reaching a diameter of ca. 5–8 µm, while plastids in phloem cells were smaller (ca. 3–6 µm). The mitochondria in the *M. scorpioides* pollen grains analyzed in the present study were small (0.19–0.20 µm). Similarly, Liu (2012) reported that the diameter of mitochondria in spindle-shaped generative cells of *Citrullus lanatus* cv. Jinzhongguanlong was ca. 0.16 µm, while in the newly born generative cells reached ca. 0.5 µm. In turn, mitochondria in plant cells were shown to have a length of 1–3 µm and an approximately 0.5 µm diameter (Møller et al., 2021).

The high number of ribosomes present in the *M. scorpioides* pollen grains indicates the possibility of intensive protein biosynthesis. It has been shown that the formation of polyribosomes and the synthesis of new proteins occurring in pollen grains determine pollen tube growth (Mascarenhas, 1993).

The sperm cells and the vegetative nucleus in *M. scorpioides* were located in the pollen grains semi-medially on both sides of the equatorial constriction. A similar arrangement of these structures in pollen grains was observed in *M. azorica* and *M. laxa* (Strittmatter & Galati, 2001). Similarly, the same arrangement of sperm cells and the vegetative nucleus in pollen grains were found in *Cryptantha intermedia*, a representative of Boraginaceae, which has pollen grains of a similar type (Hargrove & Simpson, 2003).

The cross-sections of the pollen sporoderm examined in this study revealed the presence of pollenkit, which was visible on the surface of the exine and between the bacula. Strittmatter and Galati (2001) also described the presence of pollenkit on the surface of pollen grains of two other *Myosotis* species. Pollenkit is produced by the anther tapetum. One of its many functions is its involvement in nutrition and pollen development (Pacini & Hesse, 2005).

The images of the analyzed pollen grains with the pollen tube prove that the germinative pores were located in the colpi in the equatorial part of the pollen grains. Therefore, the present observations confirm the findings reported by Strittmatter and Galati (2001), showing that colpi serve a functional role in pollen grain germination. The authors also found that, after release from anthers, *M. azorica* and *M. laxa* pollen grains

are ready for rapid germination. The present observations indicate a similar phenomenon in *M. scorpioides*.

Another function is played by pseudocolpi, which were arranged alternately with colpi. According to the current knowledge, pseudocolpi in heterocolpate pollen grains perform the function in harmomegathy, which allows controlling the processes of hydration and dehydration. In some Boraginaceae, including *Myosotis*, the presence of pseudocolpi determines the particular flexibility of the pollen wall (Volkova et al., 2013). Harmomegathy enables pollen grains to adapt to dispersal when they are partially dehydrated and to take up water before they emit pollen tubes. This is possible due to the elasticity of pollen walls provided by the special structure of furrows, which are flexible and facilitate changes in volume (Božič & Šiber, 2020; Katifori et al., 2010; Pacini & Franchi, 1999; Taia, 2022).

5. Conclusions

The plasmodial tapetum in *M. scorpioides* anthers is active in pollen loculi from the microspore phase to the late phase of pollen grain development, as the periplasmodium formed from the protoplasts of tapetum cells surrounds these grains in the 3-celled phase.

The orbicules covering the inner surface of the anther wall have a polygonal shape, which has not often been evidenced in plants so far.

The numerous organelles contained in pollen grains (plastids, mitochondria) are substantially smaller than those contained in plant somatic cells. The abundant distribution of ribosomes in both the central and distal parts of pollen grains indicates the possibility of intensive protein synthesis, which usually accompanies germination and pollen tube growth.

References

- Beug, H. J. (2004). *Leitfaden der Pollenbestimmung für Mitteleuropa und Angrenzende Gebiete*. Verlag Dr. Friedrich Pfeil.
- Boussardon, C., Carrie, C., & Keechet, O. (2023). Comparison of plastid proteomes points towards a higher plastidial redox turnover in vascular tissues than in mesophyll cells. *Journal of Experimental Botany*, 74(14), 4110–4124. <https://doi.org/10.1093/jxb/erad133>
- Božič, A., & Šiber, A. (2020). Mechanical design of apertures and the infolding of pollen grain. *PNAS*, 117(43), 26600–26607. <https://doi.org/10.1073/pnas.2011084>
- Çetinbaş, A., & Ünal, M. (2015). Anther ontogeny and microsporogenesis in *Helianthus annuus* L. (Compositae). *Notulae Scientia Biologicae*, 7(1), 52–56. <https://doi.org/10.15835/nsb.7.1.9474>
- Chacón, J., Luebert, F., Hilger, H. H., Ovchinnikova, S., Selvi, F., Cecchi, L., Williams, C. M., Hasenstab-Lehman, K., Sutorý, K., Simpson, M. G., & Weigend, M. (2016). The borage family (Boraginaceae s. str.): A revised infrafamilial classification based on new phylogenetic evidence, with emphasis on the placement of some enigmatic genera. *Taxon*, 65(3), 523–546. <https://doi.org/10.12705/653.6>

- Díez, M. J., & Valdés, B. (1991). Pollen morphology of the tribes Eritrichieae and Cynoglosseae (Boraginaceae) in the Iberian Peninsula and its taxonomic significance. *Botanical Journal of the Linnean Society*, 107(1), 49–66. <https://doi.org/10.1111/j.1095-8339.1991.tb00214.x>
- Esau, K. (1973). *Anatomia roślin* [Plant anatomy]. Państwowe Wydawnictwo Rolnicze i Leśne.
- Evert, R. F. (2006). *Esau's plant anatomy. Meristems, cells, and tissues of the plant body: Their structure, function, and development*. John Wiley & Sons, Inc.
- Fernando, D. D., & Cass, D. D. (1994). Plasmodial tapetum and pollen wall development in *Butomus umbellatus* (Butomaceae). *American Journal of Botany*, 81(12), 1592–1600. <https://doi.org/10.2307/2445337>
- Fletcher, N., & Tomblin, G. (2005). *The easy wildflower guide*. Duncan Petersen Publishing.
- Fujiwara, M. T., Hashimoto, H., Kazama, Y., Hirano, T., Yoshioka, Y., Aoki, S., Sato, N., Itoh, R. D., & Abe, T. (2010). Dynamic morphologies of pollen plastids visualised by vegetative-specific FtsZ1–GFP in *Arabidopsis thaliana*. *Protoplasma*, 242, 19–33. <https://doi.org/10.1007/s00709-010-0119-7>
- Furness, C. A. (2008). A review of the distribution of plasmodial and invasive tapeta in eudicots. *International Journal of Plant Sciences*, 169(2), 207–223. <https://doi.org/10.1086/524111>
- Furness, C. A., & Rudall, P. J. (2001). The tapetum in basal angiosperms: Early diversity. *International Journal of Plant Sciences*, 162(2), 375–392. <https://doi.org/10.1086/319580>
- Gabarayeva, N., Grigorjeva, V., & Polevova, S. (2011). Exine and tapetum development in *Symphytum officinale* (Boraginaceae). Exine substructure and its interpretation. *Plant Systematics and Evolution*, 296, 101–120. <https://doi.org/10.1007/s00606-011-0479-2>
- Galati, B. G. (2003). Ubisch bodies in angiosperms. In M. R. Chauhan & A. K. Pandey (Eds.), *Advances in plant reproductive biology* (pp. 1–21). Narendra Publishing House.
- Galati, B. G., Monacci, F., Gotelli, M. M., & Rosenfeldt, S. (2007). Pollen, tapetum and orbicle development in *Modiolastrum malvifolium* (Malvaceae). *Annals of Botany*, 99(4), 755–763. <https://doi.org/10.1093/aob/mcm011>
- García, C. C. (2002). An approach to the diversity of endothelial thickenings in Solanaceae. *Flora*, 197(3), 214–223. <https://doi.org/10.1078/0367-2530-00032>
- Germoplasm Resources Informe Network (GRIN). Retrieved April 26, 2024.
- Hargrove, L., & Simpson, M. G. (2003). Ultrastructure of heterocolpate pollen in *Cryptantha* (Boraginaceae). *International Journal of Plant Sciences*, 164(1), 137–151. <https://doi.org/10.1086/344548>
- Huysmans, S., El-Ghazaly, G., & Smets, E. (1998). Orbicules in angiosperms: Morphology, function, distribution, and relation with tapetum types. *The Botanical Review*, 64, 240–272. <https://doi.org/10.1007/BF02856566>
- Izmailow, R., & Biskup, A. (2003). Reproduction of *Echium vulgare* L. (Boraginaceae) at contaminated sites. *Acta Biologica Cracoviensia, series Botanica*, 45(1), 69–75.
- Johri, B. M., Ambegaokar, K. B., & Srivastava, P. S. (1992). *Comparative embryology of angiosperms* (Vol. 2). Springer.
- Katifori, E., Alben, S., Cerdas, E., Nelson, D. R., & Dumais, J. (2010). Foldable structures and the natural design of pollen grains. *PNAS*, 107(17), 7635–7639. <https://doi.org/10.1073/pnas.0911223>
- Kłosowski, S., & Kłosowski, G. (2006). *Flora Polski. Rośliny wodne i bagienne* [Flora of Poland. Aquatic and marsh plants]. Multico Oficyna Wydawnicza sp. z o.o.
- Liu, L. (2012). Ultrastructural study on dynamics of plastids and mitochondria during microgametogenesis in watermelon. *Micron*, 43(2–3), 412–417. <https://doi.org/10.1016/j.micron.2011.10.012>
- Lombardo, G., & Carraro, L. (1976). Tapetal ultrastructural changes during pollen development. III. Studies on *Gentiana acaulis*. *Caryologia*, 29(3), 345–349. <https://doi.org/10.1080/00087114.1976.10796674>
- Luebert, F., Cecchi, L., Frohlich, M. W., Gottschling, M., Williams, C. M., Hasenstab-Lehman, K. E., Hilger, H. H., Miller, J. S., Mittelbach, M., Nazaire, M., Nepi, M., Nocentini, D., Ober, D., Olmstead, R. G., Selvi, F., Simpson, M. G., Sutorý, K., Valdés, B., Walden, G. K., & Weigend, M. (2016). Familial classification of the Boraginales. *Taxon*, 65(3), 502–522. <https://doi.org/10.12705/653.5>
- Mascarenhas, J. P. (1993). Molecular mechanisms of pollen tube growth and differentiation. *The Plant Cell*, 5(10), 1303–1314. <https://doi.org/10.1105/tpc.5.10.1303>
- Meudt, H. M. (2016). Pollen morphology and its taxonomic utility in the Southern Hemisphere bracteate-prostrate forget-me-nots (*Myosotis*, Boraginaceae). *New Zealand Journal of Botany*, 54(4), 475–497. <https://doi.org/10.1080/0028825X.2016.1229343>
- Møller, I. M., Rasmussen, A. G., & Akenet, O. V. (2021). Plant mitochondria – Past, present and future. *Plant Journal*, 108(4), 912–959. <https://doi.org/10.1111/tbj.15495>
- Nicolau, M. R., Reposi, S. D., Bonasora, M. G., Zarlavsky, G. E., Galati, B. G., & Gotelli, M. M. (2024). Ultrastructural study of pollen and tapetum development in *Hydrocleys nymphoides*, *Alisma plantago-aquatica*, and *Sagittaria montevidensis* (Alismataceae). *Protoplasma*, <https://doi.org/10.1007/s00709-024-01989-0>
- Noroozi, M., Ghahremaninejad, F., Bogler, D., Witherspoon, J. M., Ryand, G. L., Miller, J. S., Riahi, M., & Cohen, J. I. (2022). Parsing a plethora of pollen: The role of pollen size and shape in the evolution of Boraginaceae. *Cladistics*, 38(2), 204–226. <https://doi.org/10.1111/cla.12488>
- Oak, M. K., Yang, S., Choi, G., & Song, J. H. (2022). Systematic palynology in Korean Piperales with special focus on its exine surface ornamentation and orbicle morphology. *Scientific Reports*, 12(1), Article 4142. <https://doi.org/10.1038/s41598-022-08105-3>
- Pacini, E. (1996). Tapetum types in the Compositae. In D. J. N. Hind & H. J. Beentje (Eds.), *Compositae: Systematics. Proceedings of the International Compositae Conference*, Kew, 1994 (pp. 21–28).
- Pacini, E., & Franchi, G. G. (1999). Pollen grain sporoderm and types of dispersal units. *Acta Societatis Botanicorum Poloniae*, 68(4), 362–366. <https://doi.org/10.5586/asbp.1999.042>
- Pacini, E., Franchi, G. G., & Hesse, M. (1985). The tapetum: Its form, function, and possible phylogeny in Embryophyta.

- Plant Systematics and Evolution*, 149, 155–185.
<https://doi.org/10.1007/BF00983304>
- Pacini, E., & Hesse, M. (2005). Pollenkitt – Its composition, forms and functions. *Flora*, 200(5), 399–415.
<https://doi.org/10.1016/j.flora.2005.02.006>
- Podbielkowski, Z., & Sudnik-Wójcikowska, B. (2003). *Słownik Roślin Użytkowych* [Dictionary of useful plants]. Multico Oficyna Wydawnicza.
- Podhajska, Z., & Rivola, M. (1992). *Wielka Encyklopedia Roślin* [The great encyclopedia of plants]. Wydawnictwo Aventinum.
- Rao, B. H., & Rao, P. S. P. (1992). Sporogenesis and gametogenesis of some Boraginaceae. *Feddes Repertorium*, 103(1–2), 35–40.
<https://doi.org/10.1002/fedr.19921030106>
- Reynolds, E. S. (1963). The use of lead citrate at high pH as an electronopaque stain in electron microscopy. *Journal of Cell Biology*, 17(1), 208–212.
- Ruggiero, F., & Bedini, G. (2018). Systematic and morphologic survey of orbicules in allergenic angiosperms. *Aerobiologia*, 34(3), 405–422.
<https://doi.org/10.1007/s10453-018-9522-x>
- Ruggiero, F., & Bedini, G. (2020). Phylogenetic and morphologic survey of orbicules in angiosperms. *Taxon*, 69(3), 543–566. <https://doi.org/10.1002/tax.12281>
- Rutkowski, L. (2006). *Klucz do oznaczania roślin naczyniowych Polski niżowej*. 2nd ed. [The key to the determination of vascular plants of lowland Polish. 2nd ed.]. Wydawnictwo Naukowe PWN.
- Stanley, R. G., & Linskens, H. F. (1974). *Pollen: Biology, biochemistry, management*. Springer-Verlag.
- Strittmatter, L. I., & Galati, B. G. (2001). Pollen development in *Myosotis azorica* and *M. laxa* (Boraginaceae). *Phytomorphology*, 51(1), 1–9.
- Taia, W. K. (2022). Pollen grain diversity and application in taxonomy and evolution. *Taekholmia*, 42(1), 27–42.
<https://doi.org/10.21608/taec.2022.258213>
- Tiwari, S. C., & Gunning, E. S. (1986). An ultrastructural, cytochemical and immunofluorescence study of postmeiotic development of plasmodial tapetum in *Tradescantia virginiana* L. and its relevance to the pathway of sporopollenin secretion. *Protoplasma*, 133, 100–114. <https://doi.org/10.1007/BF01304626>
- Vardar, F., & Yavuz, E. A. (2018). Development of male and female gametophyte in *Symphytum orientale* L. (Boraginaceae). *Notulae Scientia Biologicae*, 10(1), 124–129. <https://doi.org/10.15835/nsb10110191>
- Verstraete, B., Moon, H. K., Smets, E., & Huysmans, S. (2014). Orbicules in flowering plants: A phylogenetic perspective on their form and function. *The Botanical Review*, 80, 107–134. <https://doi.org/10.1007/s12229-014-9135-1>
- Vinckier, S., & Smets, E. (2002a). Morphological and ultrastructural diversity of orbicules in relation to evolutionary tendencies in Apocynaceae s.l. *Annals of Botany*, 90(5), 647–662.
<https://doi.org/10.1093/aob/mcf243>
- Vinckier, S., & Smets, E. (2002b). Morphology, ultrastructure and typology of orbicules in family Loganiaceae s.l. and related genera, in relation to systematics. *Review of Palaeobotany and Palynology*, 119(3–4), 161–189.
[https://doi.org/10.1016/S0034-6667\(01\)00137-3](https://doi.org/10.1016/S0034-6667(01)00137-3)
- Vinckier, S., & Smets, E. (2003). Morphological and ultrastructural diversity of orbicules in Gentianaceae. *Annals of Botany*, 92(5), 657–672.
<https://doi.org/10.1093/aob/mcgl87>
- Volkova, O. A., Severova, E. E., & Polevova, S. V. (2013). Structural basis of harmomegathy: Evidence from Boraginaceae pollen. *Plant Systematics and Evolution*, 299, 1769–1779. <https://doi.org/10.1007/s00606-013-0832-8>
- Volkova, O., Severova, E., & Polevova, S. (2017). Development of heterocolpate pollen in *Myosotis scorpioides* L. (Cynoglosseae, Boraginaceae). *Grana*, 56(5), 368–376.
<https://doi.org/10.1080/00173134.2016.1276211>
- Weigend, M., Selvi, F., Thomas, D. C., & Hilger, H. H. (2016). Boraginaceae. In J. W. Kadereit & V. Bittrich (Eds.), *Flowering plants. Eudicots. The families and genera of vascular plants* (Vol. 14, pp. 41–102). Springer International Publishing.
- Wilson, Z. A., Song, J., Taylor, B., & Yang, C. (2011). The final split: The regulation of anther dehiscence. *Journal of Experimental Botany*, 62(5), 1633–1649.
<https://doi.org/10.1093/jxb/err014>